## SPATIAL DIFFUSION MODELING IN SIMULATION OF SUBURBAN SPRAWL

# A Case Study in the Chicago Metropolitan Region

Y.Q. Wang and Yuming Wen


Department of Natural Resources Science, University of Rhode Island, Kingston, Rhode Island 02881


#### **INTRODUCTION**

Spatial diffusion phenomena occur because of the existence of a concentration gradient of information or material flow. The spatio-temporal process of suburban sprawl manifests the dynamic mechanism of spatial diffusion, the existence of concentration gradient. In this study, we developed a new methodology, spatial diffusion (SDIF) model to simulate suburban sprawl in the Chicago metropolitan region. Remote sensing, GIS and Census data were applied in the modeling process.


#### **DATA**


- Land cover data of 1972, 1985 and 1997 obtained from classification of Landsat images, and other available GIS data.
- Projection of the region's population, households and employment for the year 2020, which was made by the US Census Bureau and endorsed by the Northeastern Illinois Planning Commission (Drivers).
- Federal land, natural preserves, state conservation, open water, state parks, and state fish and wildlife areas (Barriers).
- Transportation, Federal or State policy.





#### MODELING PROCEDURES





### **DISCUSSION**

Most traditional models for urban sprawl are static, and are not efficient nor effective for simulating urban sprawl phenomena. Based on urban sprawl's dynamic characteristics, and according to the mechanism of spatial diffusion, a new modeling method for urban sprawl has been developed: spatial diffusion(SDIF) modeling. Since the SDIF model considers the spatial dynamics of urban sprawl phenomena, and also as many factors as possible, it is a feasible way to predict the trend of urban sprawl in any metropolitan region.

**ACKNOWLEDGEMENT:** This research is funded by NASA(grant NAG5-8829) NIP in Earth Science.